

# SureKill<sup>®</sup> ROACH GEL BAIT

SureKill Roach Gel Bait is a ready-to-use insecticide product to control roaches

- Kills cockroaches where they hide, breed and live
- Use indoors and outdoors adjacent to home or structure

**KILLS:** Cockroaches, including American, German and Oriental, Smokey Brown, and other species.

**STOP – READ THE LABEL BEFORE USE**  
**KEEP OUT OF REACH OF CHILDREN**

See backside for complete label, including directions for use and storage and disposal.

**Net Contents:** 1.4 oz (40 g) cartridge

<b>ACTIVE INGREDIENT:</b>	
Imidacloprid.....	2.15%
<b>OTHER INGREDIENTS:</b> .....	97.85%
<b>TOTAL</b> .....	100.00%
EPA Reg. No. 92028-3-47000 EPA Est. No.92418-ESP-1	

See backside for complete label, including directions for use and storage and disposal.

SPECIMEN LABEL

# SureKill<sup>III</sup> ROACH GEL BAIT

SureKill Roach Gel Bait is a ready-to-use insecticide product to control roaches

- Kills cockroaches where they hide, breed and live
- Use indoors and outdoors adjacent to home or structure

**KILLS:** Cockroaches, including American, German and Oriental, Smokey Brown, and other species.

**FOR USE IN AND AROUND STRUCTURES:** As a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches. Use SureKill Roach Gel Bait as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments, including: warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, apartment buildings, bakeries, beverage plants, breweries, bottling facilities, candy plants, canneries, cereal processing and manufacturing plants, dairies and dairy product processing plants, frozen food plants, schools, laboratories, computer facilities, aircraft, buses, boats/ships, trains, homes, industrial buildings, office buildings, animal boarding, kitchens, manufacturing facilities, spice plants, stores, wineries and similar structures, mausoleums, meat processing and packaging plants, meat and vegetable canneries, pet shops, and zoos. Non-food/non-feed areas include: bedrooms, living rooms, bathrooms, closets, media rooms, garage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, basements, attics, crawl spaces, mop closets, and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as specified in the Food/Feed Handling Establishments section.

If product is applied with an applicator gun, the gun should be suitable to fit 1.4 oz (40gm) cartridges.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label. Use strictly in accordance with label precautionary statements and directions.

### USE RESTRICTIONS:

- USE ONLY IN AREAS NOT EASILY ACCESSIBLE TO CHILDREN AND PETS.
- Do not treat food preparation surfaces. Keep bait gel away from foods and food contact surfaces.
- Clean up spilled food and debris before applying bait gel to dry surfaces.
- Do not apply bait gel to areas which have been recently sprayed with insecticide, and do not spray insecticide over bait gel as it may cause the bait gel to become repellent.
- Do not apply bait gel in areas where there is risk of electrical shock.

### APPLICATION DIRECTIONS:

Product may stain textiles, fabrics, and porous surfaces. Avoid applying to fabric or carpeting to prevent staining.

**Application rate for all uses:** Remove cap from tip and apply bait gel to treatment area. Replace cap after use. Application rate varies according to the species of cockroach to be controlled. Apply bait gel as rounded spots (approximately 1/8" inch or 4 mm in diameter) or in line approximately 1/8-inch wide and varying lengths according to the application table below. Each rounded drop weighs approximately 0.0014 ounce (0.04 grams).

Apply into cracks, crevices, holes and other small openings where roaches may be a problem. Place bait gel close to cockroach harborages, foraging, and feeding areas such as corners, cracks and crevices, and places where cockroaches may have access into structures. Numerous smaller bait gel placements provide more opportunities for cockroaches to feed on the bait gel than fewer larger placements.



Schematic representation of the drop size

Indoor and Outdoor Use Table: Match application rates according to the area and species of cockroach being treated.

Cockroach Species	Minimum Application Rate per square yard		Application Rate per 100 square feet		
	Spots	Oz.	Spots	Oz.	Total Bead Length (inches)
German	3	0.0047	37	0.0522	7
Oriental	5	0.0071	56	0.0789	10
American	5	0.0071	56	0.0789	10

Use the table to determine the number of spots or line length of SureKill Roach Gel Bait according to the species of cockroach. Inspect placements of bait gel every one to two weeks. Replenish bait gel based on the amount consumed or, replace dried out bait with fresh bait as needed.

**Indoor Use:** Apply bait gel directly into cracks and crevices leading into wall, ceiling or floor voids - areas where insects harbor. Apply bait gel under and/or behind equipment and appliances (such as: dishwashers, refrigerators, ice machines, washing machines), sinks and countertops; behind electrical switch plates. Apply gel bait around plumbing pipes and openings leading into wall voids next to tubs and toilets; around windows, doors, and potential points of entry.

**Food/Feed Handling Establishments:** Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. If bait gel contacts an exposed surface where food is handled, remove bait gel, wash exposed surface and rinse prior to use.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed, and facility is in operation. Apply bait gel in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases; around plumbing pipes, doors and windows; behind and under refrigerators, cabinets, sinks, stoves, and other equipment where cockroaches hide. During follow-up visits inspect bait gel placements and reapply when necessary.

### Use Restrictions in Food/Feed Handling Establishments:

- Do not apply bait gel to areas where food/feed, utensils, or processing surfaces may become contaminated.
- Do not apply bait gel to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation, and prepared food holding surfaces as bait gel may be removed by washing.
- Do not deposit bait gel onto exposed surfaces. If bait gel contacts an exposed surface, remove bait gel and wash exposed surface.
- Do not contaminate food and feedstuffs.

**Outdoor Use - adjacent to home or structure:** Cockroaches typically infest dark, warm and damp areas. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness. Priority treatment areas may be identified by visual inspection or aided by trapping techniques or spot flushing.

Place bait gel in areas adjacent to structures where roaches may nest or breed, or potential points of insect entry. Apply bait gel directly into cracks and crevices of walls, building voids, around sheds, porches, and other outdoor buildings, behind siding and molding, around windows, doors, sills, eaves and soffits, in expansion joints, and pipes where roaches may enter the building. Inspect bait placements every one to two weeks and replenish bait based on the amount consumed.

## STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Non-refillable container. Do not reuse this container. **If empty:** Offer container for recycling, if available, or place in trash if allowed by state and local regulations. **If partially filled:** Call your local solid waste agency for disposal instructions.

### WARRANTY

Read the entire Directions for Use and Warranty before using this product. By using this product, user or buyer accepts the following warranty. The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer. The seller is committed to providing high quality products. To the extent consistent with applicable law, the seller makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, the manufacturer or the seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.

For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.

Non-Emergency call: 1-800-621-8829

Manufactured For: Chem-Tech, Ltd.<sup>TM</sup>  
1006 Business Highway 5  
Pleasantville, IA 50225 U.S.A.

L7603-1121

